

Extended nonlocal chiral-quark model for the heavy-light quark systems

Seung-il Nam*

School of Physics, Korea Institute for Advanced Study (KIAS), Seoul 130-722, Republic of Korea

(Dated: August 28, 2012)

In this talk, we report the recent progress on constructing a phenomenological effective model for the heavy-light quark systems, which consist of (u, d, s, c, b) quarks, i.e. extended nonlocal chiral-quark model (ExNLChQM). We compute the heavy-meson weak-decay constants to verify the validity of the model. From the numerical results, it turns out that $(f_D, f_B, f_{D_s}, f_{B_s}) = (207.54, 208.13, 262.56, 262.39)$ MeV. These values are in relatively good agreement with experimental data and various theoretical estimations.

Keywords: Heavy-quark effective field theory, charm and bottom quarks, heavy mesons, instanton, nonlocal chiral-quark model

I. INTRODUCTION

The heavy-quark effective field theory (HQEFT) is a very powerful theoretical tool to investigate various heavy hadrons [1]. HQEFT manifests the heavy-quark symmetries: Spin and heavy-flavor symmetries [1]. In general, they are not obvious in full QCD but become relevant in the heavy-quark limit $m_Q \rightarrow \infty$. Using HQEFT, the light-quark component of the heavy meson was treated nonperturbatively, manifesting the spontaneous breakdown of chiral symmetry (SBCS), in many different methods [2–5]. In the present talk, we report the recent progress on constructing a *new* effective model for the heavy hadrons. The model is based on two ingredients in principle: HQEFT and nonlocal chiral-quark model (NLChQM) for the heavy and light quarks, respectively [6–8], resulting in the extended NLChQM (ExNLChQM). Intuitively, the heavy-meson mass can be regarded as the total sum of the various light and heavy quark masses, i.e. the effective and current ones:

$$\underbrace{1869 \text{ MeV}}_{D \text{ meson mass}} > \underbrace{350 \text{ MeV}}_{\text{light quark effective mass}} + \underbrace{1290 \text{ MeV}}_{\text{heavy quark current mass}}. \quad (1)$$

From Eq. (1), one needs additional 200 MeV mass. Hence, we assume that this additionally (*effective*) mass can be generated from the similar mechanism with the light quark, i.e. from the nontrivial contributions from the QCD vacuum. For more details, one can refer [9, 10]. This is a main idea to construct ExNLChQM. We organize the report as follows: In Section II, we discuss the formulation of ExNLChQM. Section III is devoted to the numerical results with relevant discussions. Summary and conclusion will be given in Section IV.

II. THEORETICAL FRAMEWORK

Using a generic functional integral technique, we can arrive at an effective action for the heavy-light quark systems from the effective Lagrangian density of ExNLChQM [9, 10]:

$$\mathcal{S}_{\text{eff}}^{\text{LL+HL+LH}} = -i \text{Sp} \ln \left[- \left(\frac{1}{f_H} \sqrt{M_Q} H \sqrt{M_q} \right) (iv \cdot \partial - M_Q)^{-1} \left(\frac{1}{f_H} \sqrt{M_q} \bar{H} \sqrt{M_Q} \right) \right], \quad (2)$$

where $\bar{M}_q = m_q + M_q$. The momentum-dependent effective masses for the heavy and light quarks are given by

$$M_{(q,Q)} = M_{(q,Q),0} \left[\frac{2\Lambda_{(q,Q)}^2}{2\Lambda_{(q,Q)}^2 - |i\not{p}|^2} \right]^2, \quad (3)$$

where $(\Lambda_q, \Lambda_Q) = (600, 1000)$ MeV [9, 10]. f_H denotes the weak-decay constant for the heavy meson. We determine $M_{Q,0}$, which is the effective heavy-quark mass at zero virtuality, from a simple phenomenological analysis:

$$M_H \approx [m_q + M_{q,0}]_L + [m_Q + M_{Q,0}]_H. \quad (4)$$

*E-mail: sinam@kias.re.kr




FIG. 1: (Color online) f_{D,D_s} (left) and f_{B,B_s} (right) as functions of Λ_Q at $\Lambda_q = 600$ MeV. The numerical results for the non-strange and strange mesons are shown with the solid and dot lines, respectively. The experimental data are taken from [11, 12]. Note that the experimental errors are given with the shaded areas.

From the experimental data for D and B mesons, we can estimate the values of $M_{Q,0}$ for numerical calculations with $M_{q,0} \approx 350$ MeV [7]:

$$\begin{aligned} M_D &= 1869.57 \text{ MeV} \approx 1625.0 \text{ MeV} + M_{Q,0} \rightarrow M_{Q,0} \approx 244.57 \text{ MeV} \\ M_B &= 5279.17 \text{ MeV} \approx 5025.0 \text{ MeV} + M_{Q,0} \rightarrow M_{Q,0} \approx 254.17 \text{ MeV}. \end{aligned} \quad (5)$$

The heavy-meson weak-decay constant, f_H is defined by the following:

$$\langle 0 | \bar{q}(x) \gamma_\mu (1 - \gamma_5) \tau^a Q_v(x) | H^b(p) \rangle = 2ip_\mu f_H \delta^{ab}. \quad (6)$$

The matrix element in the left-hand-side of Eq. (6) can be easily derived from the functional derivative of the effective chiral action in Eq. (2).

III. NUMERICAL RESULTS

In Fig. 1, we depict the numerical results for f_H as functions of Λ_Q by fixing $\Lambda_q = 600$ MeV. The horizontal lines indicate the global fit data with errors (shaded areas). To reproduce the data, we fix $\Lambda_Q = 1$ GeV for every heavy-quark flavors as shown in the figure and mentioned in Sect. 2. From those Λ parameters, we have $(f_D, f_B, f_{D_s}, f_{B_s}) = (207.54, 208.13, 262.56, 262.39)$ MeV. Hence, we have the following tendency from the present model:

$$(f_D \approx f_B \approx 210 \text{ MeV}) < (f_{D_s} \approx f_{B_s} \approx 260 \text{ MeV}). \quad (7)$$

These computed values are well compatible with other theoretical estimations including lattice QCD simulations. Detailed comparisons are given in Fig. 2.

IV. SUMMARY AND CONCLUSION

In the present work, we have investigated the weak-decay constants for the heavy PS-mesons consisting of the (u, d, s, c, b) flavors, i.e. f_{D,B,D_s,B_s} , using ExNLChQM. We list two important observations in the present work:

- We employ a phenomenological correction factor for the inclusion of the strange quark, compensating the $1/N_c$ corrections. By doing that, we obtain $f_{D,D_s,B,B_s} = (207.53, 262.56, 208.13, 262.39)$ MeV, which are qualitatively compatible with available experimental and theoretical values.
- It is justified that the vacuum structure due to the heavy quarks is modified in comparison to that for the light quarks by seeing that $\Lambda_q \neq \Lambda_Q$. This tendency also signals the breakdown of the SU(5) flavor symmetry at the quark level, in addition to $m_q \neq m_Q$.




FIG. 2: (Color online) Theoretical results for f_{D,D_s,B,B_s} from the present calculations, clover-improved quenched LQCD [13], field-correlator method (FC) [3], QCD sum rule [14], light-front quark model (LQM) [15], Bethe-Salpeter method (BS) [16], relativistic quark model (RQM) [2], and light-cone wave function (LC) [4].

Acknowledgments

The present report is prepared for the proceedings of the 20th International IUPAP Conference on Few-Body Problems in Physics (FB20), 20 ~ 25 August 2012, Fukuoka, Japan. The author would like to thank A. Hosaka for fruitful discussions on this subject.

-
- [1] H. Georgi, Phys. Lett. **B240**, 447-450 (1990).
 - [2] D. Ebert, R. N. Faustov and V. O. Galkin, Phys. Lett. B **635**, 93 (2006).
 - [3] A. M. Badalian, B. L. G. Bakker and Yu. A. Simonov, Phys. Rev. D **75**, 116001 (2007).
 - [4] C. W. Hwang, Phys. Rev. D **81**, 114024 (2010).
 - [5] L. S. Geng, M. Altenbuchinger, W. Weise, Phys. Lett. **B696**, 390-395 (2011).
 - [6] M. Praszalowicz and A. Rostworowski, Phys. Rev. D **64**, 074003 (2001).
 - [7] S. i. Nam, H. -Ch. Kim, A. Hosaka, M. M. Musakhanov, Phys. Rev. **D74**, 014019 (2006).
 - [8] A. E. Dorokhov, JETP Lett. **92**, 707 (2010).
 - [9] S. i. Nam, Phys. Rev. D **85**, 034019 (2012).
 - [10] S. i. Nam, Mod. Phys. Lett. A **26**, 1250153 (2012).
 - [11] J. L. Rosner, S. Stone, [arXiv:0802.1043 [hep-ex]].
 - [12] J. L. Rosner, S. Stone, [arXiv:1002.1655 [hep-ex]].
 - [13] D. Becirevic *et al*, Phys. Rev. D **60**, 074501 (1999).
 - [14] A. A. Penin and M. Steinhauser, Phys. Rev. D **65**, 054006 (2002).
 - [15] H. M. -Choi, Phys. Rev. D **75**, 073016 (2007).
 - [16] G. -Chvetic, C. S. Kim, G. L. Wang and W. Namgung, Phys. Lett. B **596**, 84 (2004).